



PEST CONTROL STUDY MOLT-X[®]:

RESULTS OF TESTING

Treated with 1 application each of 8, 16 and 32 oz./100 gallons, unless otherwise noted (*).

BOTANICAL NAME	PLANT SPECIES (VARIETY)	FOLIAR PHYTOTOXICITY (YES/NO)	FLOWER PHYTOTOXICITY (YES/NO)
Ageratum houstonianum	Ageratum (floss flower)	N	N
Baptisia australis*	Blue false indigo	N	**
Begonia	Begonia	N	N
Begonia semperflorens	Wax leaf Begonia	N	Y
Calibrachoa hybrid*	Calibrachoa	N	Y
Catharanthus roseus	Vinca (periwinkle)	N	Y
Ceratostigma plumbaginoides*	Plumbago	N	N
Coreopsis verticillata	Coreopsis (Moonbeam)	N	N
Dianthus gratianopolitanus	Dianthus (Firewitch)	N	N
Eriathus ravennae	Hardy Papas Grass	Y	**
Geranium*	Hardy geranium (Rosanne)	N	Y
Gerbera jamesonii	Gerbera Daisy	N	N
Hypericum calycinum	St. John's Wort	N	N
Iberis sepevires	Iberis (Purity white)	N	N
Impatiens × hybrida	New Guinea Impatiens	N	**
Impatiens walleriana	Impatiens (mixed)	N	N
Pelargonium	Geranium (Deep Pink)	N	N
Salvia nemorosa	Salvia (May Night)	N	**
Salvia nemorosa	Salvia (red)	N	N
Sedum spectabile	Sedum (Neon)	Y	**
	Tomato (Cherokee Purple)	N	**
Stokesia laevis	Stokes Aster (Color Wheel)	N	N
Tagetes erecta	Marigolds (African)	N	N
Torenia fournieri	Torenia	N	N
Verbena Canadensis*	Verbena (Snow Flurry)	N	N
Zinnia	Zinnia	N	N

*Treated with 1 application at 8, 10 and 12 oz / 100 gallons.

**No bloom present at time of testing.

NOTE: Use caution when making Molt-X applications to open blooms, especially on varieties known to be sensitive. It is recommended that a small group of plants be tested for effects on open blooms before making applications to the whole crop. Always read and follow label directions.