

CROP GUIDE VEGETABLE TRANSPLANTS & HERBS

	CROP STAGE			
PEST/DEFICIENCY	SEED/PROPAGATION	SEEDLING (3-5 LEAVES)	VEGETATIVE	
Soilborne Diseases	RootShield® <i>PLUS†</i> WP, RootShield PLUS+ <i>G</i>	RootShield <i>PLUS⁺</i> WP		
Powdery Mildew, Downy Mildew		MilStop® SP + CEASE®, PERpose Plus		
Botrytis	PERpose Plus®	BotryStop® WP, MilStop SP + CEASE		
Bacterial Leaf Spot		PERpose Plus, CEASE		
Aphids, Whiteflies, and Thrips	EpiShield™			
Fungus Gnats, Shore Flies	NemaShield®			
Mites	EpiShield SuffOil-X ⁴ , EpiShield		, EpiShield	

Efficient nutrient use and stress reduction	ON-Gard®		
Calcium (Ca) deficiency	ON-Gard CALCIUM		



ABOVE AND BEYOND IS WHERE WE BEGIN

CROP GUIDE VEGETABLE TRANSPLANTS & HERBS

		PRODUCT	REI, PHI	OMRI	PEST/DEFICIENCY	RECOMMENDED RATE	TYPE OF APPLICATION
DISEASES		RootShield <i>PLUS</i> + WP²	4-hour 0-day	Yes	Damping Off, Pythium, Fusarium, Phytophthora, Rhizoctonia	Drench: 4-8 oz/100 gal	Drench at seeding.
		RootShield <i>PLUS</i> ⁺ G ²	4-hour 0-day	Yes		1.5-3 lb/cu yd	Incorporate in potting media.
		MilStop SP	1-hour O-day	Yes	Powdery Mildew, Downy Mildew, <i>Botryti</i> s	1.25-5.0 lb/100 gal	Foliar application every 7-14 days. Tank mix rates: CEASE 4-6 qt/100 gal + MilStop SP 1.25-3.0 lb/100 gal.
		CEASE	4-hour 0-day	Yes	Bacterial Leaf Spot, Powdery Mildew, Downy Mildew, <i>Botrytis</i>	4-6 qt/100 gal	Foliar application every 3–7 days. Safe for use with young seedlings.
		PERpose Plus¹	0-1-hour 0-day	Yes		1/3–1 fl oz/gal Mist: 1:1000–1:5000	1/3 fl oz/gal preventative foliar application weekly; 1 fl oz/gal curative application. Mist: Apply 1:1000 or 7-10 days at seeding, continue 1:5000 into propagation.
TES		BotryStop WP	4-hour 0-day	Yes	Botrytis	2-4 lb/100 gal	Foliar application every 7-10 days.
		Molt-X ²	4-hour 0-day	Yes	Aphids, Whiteflies, Thrips	8–10 fl oz/100 gal	Foliar spray to glisten on a 7-10-day interval or as needed.
IW / S.		SuffOil-X ⁴	4-hour 0-day	Yes	Aphids, Whiteflies, Thrips, Mites	1-2 gal/100 gal	Foliar application every 7 days.
INSECT		EpiShield	N/A	No	Mites, Aphids, Thrips, Whiteflies, Mealybugs	Foliar spray 9-12 fl oz/100 gal Dip rate 2.7 ml/gal	Foliar spray to glisten on 5-7-day interval or as needed. Do not exceed 9 fl oz rate in flower. Dip unrooted cuttings for 10-30 seconds while moving plants around in solution.
		NemaShield	N/A	N/A	Fungus Gnats, Shore Flies	100 million/1100-3400 ft²	Apply as a drench to the substrate as needed.
		Molt-X ²	4-hour 0-day	Yes		8 fl oz/100 gal	Apply as a drench.
NUTRITION		ON-Gard (5-0-0)³	N/A Yes	Nutrition supplement and stress reduction	Drench, sprench, or foliar spray: 1 qt/100 gal	Drench applications at seeding and weekly foliar applications. Fog rates: 6.4-11.2 fl oz/3 L water.	
	***	ON-Gard CALCIUM³ (1-0-0 + 5% Ca)		Calcium (Ca) deficiency	16 fl oz/100 gal	Weekly foliar spray to prevent Ca deficiency.	

Please refer to product labels for complete application instructions. Always read and follow label directions.

'Not available for use in CA; 'Not available for use in AK; 'Not labeled for use in MT, ND, SD, WY. 4SuffOil-X not labeled for use on herbs

Please refer to product labels ${\bf bioworksinc.com}$ for fogging application details and rates.



BIOWORKS, INC.

100 Rawson Road • Suite 205

Victor, NY 14564

bioworksinc.com • 800.877.9443