

CROP GUIDE GREENHOUSE TOMATOES

	CROP STAGE					
PEST/DEFICIENCY	SEED	SEEDLING (3-5 LEAVES)	VEGETATIVE STAGE	FRUIT PRODUCTION STAGE		
Soilborne Diseases	RootShield® <i>PLU</i> S⁺ WP					
Powdery Mildew		CEASE®, MilStop® SP	MilStop SP + CEASE, PERpose Plus®			
Botrytis		CEASE .	BotryStop® WP,	MilStop + CEASE		
White Mold			CEASE			
Bacterial Spot, Bacterial Speck, Early Blight			CEASE, PE	Rpose Plus		
Aphids, Whiteflies, Thrips, and Tomato Psyllid		Molt-X®, SuffOil-X®, EpiShield™				
Lepidopterans			Molt-X			
Tarnished Plant Bugs			MOTE-X			
Fungus Gnats, Shore Flies	NemaShield, Molt-X					
Mites		SuffOil-X, EpiShield				
Efficient nutrient use and stress reduction	$ON ext{-}Gard^{\otimes}$					
Calcium (Ca) deficiency	ON-Gard CALCIUM					

ABOVE AND BEYOND IS WHERE WE BEGIN

CROP GUIDE GREENHOUSE TOMATOES

		PRODUCT	REI, PHI	OMRI	PEST/DEFICIENCY	RECOMMENDED RATE	TYPE OF APPLICATION	
INSECTS / MITES DISEASES		RootShield <i>PLUS</i> ⁺ WP ²	4-hour 0-day	Yes	Pythium, Fusarium, Phytophthora, Rhizoctonia, Damping Off	Drench: 4-8 oz/100 gal	Drench at seeding and transplant. Reapply at 4–6-week intervals.	
		MilStop SP	1-hour 0-day	Yes	– Powdery Mildew, Botrytis	1.25–5.0 lb/100 gal	Foliar application every 7–14 days. Tank mix rates: CEASE 4–6 qt/100 gal + MilStop SP 1.25–3 lb/100 gal.	
		CEASE	4-hour 0-day	Yes		4-6 qt/100 gal	Foliar application every 3-7 days.	
		PERpose Plus¹	0-1-hour 0-day	Yes		1/3–1 fl oz/gal	1/3 fl oz/gal preventative foliar application weekly; 1 fl oz/gal curative application.	
		BotryStop WP ⁴	4-hour 0-day	Yes	Botrytis	2-4 lb/100 gal	Foliar application every 7–10 days.	
	*	Molt-X²	4-hour O-day	Yes	Whiteflies, Aphids, Thrips, Psyllids, Lepidopterans, Tarnished Plant Bug	8-10 fl oz/100 gal	Foliar spray to glisten on a 7-10-day interval or as needed.	
		SuffOil-X	4-hour 0-day	Yes	Mites, Whiteflies, Aphids, Thrips, Tomato Psyllid	1-2 gal/100 gal	Foliar application every 7 days. Suppresses powdery mildew.	
		EpiShield	N/A	No	Mites, Aphids, Thrips, Whiteflies	9-12 fl oz/100 gal	Foliar spray to glisten on 5-7-day interval or as needed.	
		NemaShield	N/A	N/A	Fungus Gnats, Shore Flies	100 million/1100-3400 ft²	Apply as a drench to the substrate as needed.	
		Molt-X²	4-hour 0-day	Yes		8 fl oz/100 gal	Apply as a drench.	
NUTRITION	***	ON-Gard (5-0-0)³	N/A	Yes	Nutrition supplement and stress reduction	1 qt-2 gal/100 gal	Weekly foliar spray: 1 qt in seedling stage to 2 gal in vegetative & fruiting stages. Fog rates: 6.4-11.2 fl oz/3 L water.	
		ON-Gard CALCIUM³ (1-0-0 + 5% Ca)			Calcium (Ca) deficiency	8-16 fl oz/100 gal	Weekly foliar spray: 8 fl oz in seedling stage to 16 fl oz in vegetative & fruiting stages.	

Please refer to product labels for complete application instructions. Always read and follow label directions.

'Not available for use in CA; 'Not available for use in AK; 'Not labeled for use in MT, ND, SD, WY. 'BotryStop WP not labeled for use on tomatoes in CA

Please refer to product labels ${\bf bioworksinc.com}$ for fogging application details and rates.



BIOWORKS, INC.

100 Rawson Road • Suite 205

Victor, NY 14564

bioworksinc.com • 800.877.9443