



APPLICATION/PRODUCTION: BOTRYSTOP® WP PEONIES

BotryStop WP (*Ulocladium oudemansii* U3 Strain), a living micro-organism, provides protection against *Botrytis* infection by colonizing senescing tissue and preventing *Botrytis* establishment.

BEST PRACTICES FOR PEONIES

For successful use of BotryStop WP, the following practices should be carried out:

- + Ensure that your sprayer is correctly calibrated
- + Ensure healthy crop hygiene management by removing crop debris. This reduces the sites for *Botrytis* to over-winter
- + Time BotryStop WP applications to ensure that *Botrytis* is out-competed early
- + Use the correct surfactant for the volume of water per acre
- + Use at least 50 gallons of water or more per acre. Higher water volumes improve coverage (avoid runoff)
- + Mix with ONLY ONE tank mix partner*

*Refer to BioWorks Product Compatibility Tool at bioworksinc.com.

APPLICATION NOTES

- + BotryStop WP should be applied every 7-14 days from shoot emergence to bud break
- + Apply BotryStop WP at the rate of 3 lb/acre
- + Store BotryStop WP in a cool, dry place at or below 68° F, away from direct sunlight. Store in original packaging, with excess air removed, and bag tightly closed, away from non-compatible fungicides*
- + Add the wettable powder directly to a clean, well-rinsed spray tank with agitation running; no pre-mixing is necessary
- + For optimal performance, add a suitable surfactant to the spray tank to assist product wetting/spreading* (Note: For certified organic grapes, use an approved organic adjuvant)
- + BotryStop WP is most effective when applied at high water volumes (50 gal/acre or greater)
- + BotryStop WP can be tank mixed, but only with ONE other compatible tank mix partner.* DO NOT tank mix with two or more fungicides (e.g. Do not mix BotryStop WP with copper and sulfur)
- + When tank mixing BotryStop WP with another compatible fungicide use the manufacturer's recommended rate. Mixtures are best applied at high volume

*Refer to compatibility table at bioworksinc.com



APPLICATION/PRODUCTION BOTRYSTOP® WP: PEONIES

APPLICATION NOTES (CONTINUED)

- + Once mixed, the product should be used immediately
- + To minimize residue on open flowers, use small spray droplet sizes and spray to wet, not to runoff
- + For improved results, make applications in the early morning or late evening when temperatures are cooler and dew may be on leaves. Avoid application under hot, dry conditions, such as midday
- + Use a well calibrated/set up sprayer to ensure the entire plant is well covered during spraying
- + Maintain tank agitation until spraying is complete
- + Rinse out the tank, spray lines and nozzles after application

Refer to product labels for complete application details. Additional technical information is available at bioworksinc.com or from your BioWorks technical sales representative. BotryStop® WP is a trademark of BioWorks, Inc. Always read and follow label directions.

GOING ABOVE AND BEYOND

At BioWorks, we not only help eradicate harmful diseases and pests that threaten your crops, but we design and support integrated plant health management programs. And because we know how quickly some diseases can work, we respond to any questions you might have in just four hours. So you can get back to the business of successful growing.



BIOWORKS

100 Rawson Road • Suite 205

Victor, NY 14564

expert@bioworksinc.com

bioworksinc.com • 800.877.9443