

CROP GUIDE GREENHOUSE PEPPERS

		CROP STAGE					
	PEST/DEFICIENCY	SEED	SEEDLING (3-5 LEAVES)	VEGETATIVE STAGE	FRUIT PRODUCTION STAGE		
	Soilborne Diseases	RootShield® <i>PLU</i> S⁺ WP					
	Powdery Mildew		CEASE®, MilStop® SP	MilStop SP + CEASE, PERpose Plus®			
	Botrytis		CEASE	BotryStop® WP, M	ilStop SP + CEASE		
	White Mold		CLASE	CEASE, PERpose Plus			
	Fusarium Fruit Rot, Bacterial Soft Fruit Rot			CEASE, PERpose Plus			
	Aphids, Whiteflies, Thrips		Molt-X®, SuffOil-X®, EpiShield™				
	Lepidopterans						
	Pepper Weevils	Molt-X					
	Tarnished Plant Bugs						
	Fungus Gnats, Shore Flies	NemaShield®, Molt-X					
	Spider Mites, Broad Mites		SuffOil-X, EpiShield				
Efficient nutrient use and stress reduction ON-Gard®				Gard®			
	Calcium (Ca) deficiency				ON-Gard CALCIUM		

ABOVE AND BEYOND IS WHERE WE BEGIN

CROP GUIDE GREENHOUSE PEPPERS

		PRODUCT	REI, PHI	OMRI	PEST/DEFICIENCY	RECOMMENDED RATE	TYPE OF APPLICATION
SEASES	**	RootShield <i>PLUS</i> + WP²	4-hour 0-day	Yes	Pythium, Fusarium, Phytophthora, Rhizoctonia, Damping Off	Drench: 4-8 oz/100 gal	Drench at seeding and transplant; reapply at 4–6 week intervals.
D		MilStop SP	1-hour 0-day	Yes	Powdery Mildew, Botrytis	1.25–5.0 lb/100 gal	Foliar application every week. Tank mix rates: CEASE 4–6 qt/100 gal + MilStop SP 1.25–3 lb/100 gal.
		CEASE	4-hour 0-day	Yes	Powdery Mildew, Botrytis, White Mold (Sclerotinia)	4-6 qt/100 gal	Foliar application every 7–14 days.
		PERpose Plus¹	0-1-hour 0-day	Yes	Powdery Mildew, White Mold (S <i>clerotinia</i>)	1/3-1 fl oz/gal	1/3 fl oz/gal preventative foliar application weekly.
		BotryStop WP ⁴	4-hour 0-day	Yes	Botrytis	2-4 lb/100 gal	Foliar application every 7–10 days.
S / MITES		Molt-X²	4-hour 0-day	Yes	Aphids, Whiteflies, Thrips, Lepidopterans, Pepper Weevil, Tarnished Plant Bugs.	8-10 fl oz/100 gal	Foliar spray to glisten on a 7-10-day interval or as needed.
SECT		SuffOil-X	4-hour 0-day	Yes	Mites, Aphids, Whiteflies, Thrips	1-2 gal/100 gal	Foliar application every 7 days. Suppresses powdery mildew.
		EpiShield	N/A	No	Mites, Aphids, Thrips, Whiteflies, Mealybugs	9-12 fl oz/100 gal	Foliar spray to glisten on 5-7-day interval or as needed.
		NemaShield	N/A	N/A	Fungus Gnats, Shore Flies	100 million/1100-3400 ft2	Apply as a drench to the substrate as needed.
		Molt-X²	4-hour 0-day	Yes		8 fl oz/100 gal	Apply as a drench.
NUTRITION	─	ON-Gard (5-0-0)³	N/A	N/A Yes	Nutrition supplement and stress reduction	1 qt-2 gal/100 gal	Weekly foliar spray: 1 qt in seedling stage to 2 gal in vegetative & fruiting stages. Fog rates: 6.4-11.2 fl oz/3 L water.
FUN		ON-Gard CALCIUM³ (1-0-0 + 5% Ca)			Calcium (Ca) deficiency	8-16 fl oz/100 gal	Weekly foliar spray: 8 fl oz in seedling stage to 16 fl oz in vegetative & fruiting stages.

Please refer to product labels for complete application instructions. Always read and follow label directions.

'Not available for use in CA; aNot available for use in AK; aNot labeled for use in MT, ND, SD, WY. BotryStop WP not labeled for use on pepper in CA

Please refer to product labels ${\bf bioworksinc.com}$ for fogging application details and rates.



BIOWORKS, INC.

100 Rawson Road • Suite 205

Victor, NY 14564

bioworksinc.com • 800.877.9443