

## **CROP GUIDE POINSETTIAS**

NUTRITION

		CROP STAGE					
	PEST/DEFICIENCY	PRE-PLANT (STOCK PLANTS)	CUTTINGS (PROPAGATION)	VEGETATIVE	BRACT FORMATION		
	Soilborne Diseases	RootShield® <i>PLUS</i> <sup>+</sup> WP, RootShield <i>PLUS</i> <sup>+</sup> G					
	Powdery Mildew	CEASE®, MilStop® SP, PERpose Plus®		MilStop SP + CEASE, PERpose Plus			
	Poinsettia Scab or Spot Anthracnose						
	Bacterial Leaf Spot	PERpose Plus, CEASE, BotryStop® WP					
	Botrytis	CEASE, BotryStop WP, MilStop SP, PERpose Plus	BotryStop WP, CEASE, MilStop SP, PERpose Plus	BotryStop WP, MilStop SP + CEASE			
	Aphids, Whiteflies, Thrips	Molt-X®, SuffOil-X®, EpiShield®, PRINCIPLE™ WP					
	Mealybugs	Molt-X, SuffOil-X, EpiShield					
	Fungus Gnats, Shore Flies	NemaShield, Molt-X					
	Mites	SuffOil-X, EpiShield					
	Efficient nutrient use and stress reduction	ON-Gard®					



	PRODUCT	REI, PHI	OMRI	PEST/DEFICIENCY	RECOMMENDED RATE	TYPE OF APPLICATION	
SES	RootShield <i>PLUS</i> <sup>+</sup> WP <sup>2</sup>	4-hour 0-day	Yes	Damping Off, Pythium, Fusarium, Phytophthora,  Dip (URC): 1.5 oz/5 gal		Drench at sticking or transplant, or Dip unrooted cuttings (URC) prior to sticking. Reapply at 10 weeks.	
DISE A	RootShield <i>PLUS</i> <sup>+</sup> G <sup>2</sup>	4-hour 0-day	Yes	Rhizoctonia	2 lb/cu yd	Incorporate in potting media.	
	CEASE	4-hour 0-day	Yes	Powdery Mildew, Poinsettia Scab, Bacterial Leaf Spot	4-6 qt/100 gal	Foliar application every 3–7 days.	
	MilStop SP	1-hour 0-day	Yes	Powdery Mildew, Poinsettia Scab	1.25–5.0 lb/100 gal	Foliar application every 7-14 days. At highest rate, apply every 14 days.  Tank mix rates: CEASE + MilStop SP  1/3 fl oz/gal preventative application weekly;  1 fl oz/gal curative application.  Mist in propagation: Apply 1:1000 for first week;  1:5000 for remainder of propagation.	
	PERpose Plus	0-1-hour 0-day	Yes	Powdery Mildew, Poinsettia Scab (Spot Anthracnose), Bacterial Leaf Spot	1/3-1fl oz/gal		
	BotryStop WP	4-hour 0-day	Yes	Botrytis, Bacterial Leaf Spot¹	2-4 lb/100 gal	Foliar application every 7-10 days.	
ITES	PRINCIPLE WP1	4-hour 0-day	Yes	Whiteflies, Aphids,	0.5-2 lb/100 gal Dip rate: 0.25-0.5 oz/gal	Foliar application every 5-10 days. For thrips use 1-2 lb/100 gal.	
S. S.	Molt-X <sup>2</sup>	4-hour 0-day	Yes	Thrips, Mealybugs	8–10 fl oz/100 gal	Foliar spray to glisten on a 7-10-day interval or as needed.	
INSECT	SuffOil-X	4-hour 0-day	Yes	Mites, Aphids, Thrips, Whiteflies, Mealybugs	1-2 gal/100 gal	Foliar application every 7 days. Suppresses Powdery Mildew.	
	EpiShield	N/A	No	Mites, Aphids, Thrips, Whiteflies, Mealybugs	Foliar spray 9-12 fl oz/100 gal Dip rate 2.7 ml/gal	Foliar spray to glisten on 5-7-day interval or as needed Do not exceed 9 fl oz rate in flower. Dip unrooted cuttin for 10-30 seconds while moving plants around in solutio	
	NemaShield	N/A	N/A	Fungus Gnats, Shore	100 million/1100-3400 ft²	Apply as a drench to the substrate as needed.	
	Molt-X <sup>2</sup> 4-hour Yes		Flies	8 fl oz/100 gal	Apply as a drench.		
NUTRITION	ON-Gard (5-0-0)³	N/A	Yes	Nutrition supplement and stress reduction	Foliar Spray/Drench/Dip: 1 qt/100 gal	Pre-plant dip or drench at planting. Re-apply as a drench at 2-week intervals.	

Please refer to product labels for complete application instructions. Always read and follow label directions. 'Not available for use in CA; 'Not available for use in AK; 'Not labeled for use in MT, ND, SD, WY

Please refer to product labels **bioworksinc.com** for fogging application details and rates.

©2024 BioWorks, Inc. The following are registered trademarks of BioWorks, Inc.: RootShield, CEASE, MilStop SP, PERpose Plus, BotryStop, Molt-X, NemaShield, SuffOil-X, EpiShield, and ON-Gard. PRINCIPLE is a trademark of BioWorks, Inc.



BIOWORKS, INC.

100 Rawson Road • Suite 205

Victor, NY 14564

bioworksinc.com • 800.877.9443