

We deliver a full line of biological solutions to control pests, prevent disease and improve plant health. Our products help you grow all season and produce the highest quality poinsettias.



**BioWorks**  
A MEMBER OF BIOBEST GROUP

# CROP GUIDE POINSETTIAS

DISEASES



INSECTS / MITES



NUTRITION



PEST/DEFICIENCY	CROP STAGE			
	PRE-PLANT (STOCK PLANTS)	CUTTINGS (PROPAGATION)	VEGETATIVE	BRACT FORMATION
Soilborne Diseases	RootShield® PLUS <sup>+</sup> WP, RootShield PLUS <sup>+</sup> G			
Powdery Mildew	CEASE®, MilStop® SP, PERpose Plus®		MilStop SP + CEASE, PERpose Plus	
Poinsettia Scab or Spot Anthracnose				
Bacterial Leaf Spot	PERpose Plus, CEASE, BotryStop® WP			
Botrytis	CEASE, MilStop SP, PERpose Plus	BotryStop WP, CEASE, MilStop SP, PERpose Plus	BotryStop WP, MilStop SP + CEASE	
Aphids, Whiteflies, Thrips, Mealybugs	Molt-X®, SuffOil-X®, EpiShield™			
Fungus Gnats, Shore Flies	NemaShield, Molt-X			
Mites	SuffOil-X, EpiShield			
Efficient nutrient use and stress reduction	ON-Gard®			
Calcium (Ca) deficiency				ON-Gard CALCIUM



ABOVE AND BEYOND IS WHERE WE BEGIN

	PRODUCT	REI, PHI	OMRI	PEST/DEFICIENCY	RECOMMENDED RATE	TYPE OF APPLICATION
DISEASES	RootShield PLUS <sup>+</sup> WP <sup>2</sup>	4-hour 0-day	Yes	Damping Off, Pythium, Fusarium, Phytophthora, Rhizoctonia	Drench: 6 oz/100 gal Dip (URC): 1.5 oz/5 gal	Drench at sticking or transplant, or Dip unrooted cuttings (URC) prior to sticking. Reapply at 10 weeks.
	RootShield PLUS <sup>+</sup> G <sup>2</sup>	4-hour 0-day	Yes		2 lb/cu yd	Incorporate in potting media.
	CEASE	4-hour 0-day	Yes	Powdery Mildew, Poinsettia Scab (Spot Anthracnose), Bacterial Leaf Spot	4–6 qt/100 gal	Foliar application every 3–7 days.
	MilStop SP	1-hour 0-day	Yes	Powdery Mildew, Poinsettia Scab (Spot Anthracnose)	1.25–5.0 lb/100 gal	Foliar application every 7–14 days. At highest rate, apply every 14 days. Tank mix rates: CEASE 4–6qt/100gal + MilStop SP 1.25–3 lb/100 gal.
	PERpose Plus <sup>1</sup>	0–1-hour 0-day	Yes	Powdery Mildew, Poinsettia Scab (Spot Anthracnose), Bacterial Leaf Spot	1/3–1fl oz/gal	1/3 fl oz/gal preventative application weekly; 1 fl oz/gal curative application. Mist in propagation: Apply 1:1000 for first week; 1:5000 for remainder of propagation.
	BotryStop WP	4-hour 0-day	Yes	Botrytis, Bacterial Leaf Spot <sup>1</sup>	2–4 lb/100 gal	Foliar application every 7–10 days.
INSECTS / MITES	Molt-X <sup>2</sup>	4-hour	Yes	Whiteflies, Aphids, Thrips, Mealybugs	8–10 fl oz/100 gal	Foliar spray to glisten on a 7–10-day interval or as needed.
	SuffOil-X	4-hour	Yes	Mites, Aphids, Thrips, Whiteflies, Mealybugs	1–2 gal/100 gal	Foliar application every 7 days. Suppresses Powdery Mildew.
	EpiShield	N/A	No	Mites, Aphids, Thrips, Whiteflies, Mealybugs	Foliar spray 9–12 fl oz/100 gal Dip rate 2.7 ml/gal	Foliar spray to glisten on 5–7-day interval or as needed. Do not exceed 9 fl oz rate in flower. Dip unrooted cuttings for 10–30 seconds while moving plants around in solution.
	NemaShield	N/A	N/A	Fungus Gnats, Shore Flies	100 million/ 1100–3400 ft <sup>2</sup>	Apply as a drench to the substrate as needed.
	Molt-X <sup>2</sup>	4-hour	Yes		8 fl oz/100 gal	Apply as a drench.
NUTRITION	ON-Gard (5-0-0) <sup>3</sup>	N/A	Yes	Nutrition supplement and stress reduction	Foliar Spray/Drench/Dip: 1 qt/100 gal	Pre-plant dip or drench at planting. Re-apply as a drench at 2-week intervals. Low Volume (Fogging): 6.4–11.2 fl oz/ 3 L water.
	ON-Gard CALCIUM <sup>3</sup> (1-0-0 + 5% Ca)			Calcium (Ca) deficiency	16 fl oz/100 gal	Apply as a foliar spray at 2-week intervals.

Please refer to product labels for complete application instructions. Always read and follow label directions.

<sup>1</sup>Not available for use in CA; <sup>2</sup>Not available for use in AK; <sup>3</sup>Not labeled for use in MT, ND, SD, WY

Please refer to product labels [bioworksinc.com](http://bioworksinc.com) for fogging application details and rates.



**BioWorks**<sup>®</sup>

A MEMBER OF BIOBEST GROUP

**BIOWORKS, INC.**

100 Rawson Road • Suite 205

Victor, NY 14564

bioworksinc.com • 800.877.9443